



Contents lists available at ScienceDirect

## Placenta

journal homepage: [www.elsevier.com/locate/placenta](http://www.elsevier.com/locate/placenta)

## Mitochondrial – Endoplasmic reticulum interactions in the trophoblast: Stress and senescence

G.J. Burton<sup>\*</sup>, H.W. Yung, A.J. Murray

Centre for Trophoblast Research and Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge CB2 3EG, UK

## ARTICLE INFO

## Article history:

Received 2 February 2016

Received in revised form

30 March 2016

Accepted 1 April 2016

## Keywords:

Mitochondria

Endoplasmic reticulum

Oxidative stress

Unfolded protein response

## ABSTRACT

Placental stress has been implicated in the pathophysiology of complications of pregnancy, including growth restriction and pre-eclampsia. Initially, attention focused on oxidative stress, but recently mitochondrial and endoplasmic reticulum stress have been identified. Complex molecular interactions exist among these different forms of stress, making it unlikely that any occurs in isolation. In part, this is due to close physiological connections between the two organelles principally involved, mitochondria and the endoplasmic reticulum (ER), mediated through  $Ca^{2+}$  signalling. Here, we review the involvement of the mitochondria-ER unit in the generation of stress within the trophoblast, and consider consequences for obstetric outcome. Mild stress may induce adaptive responses, including upregulation of antioxidant defences and autophagy, while moderate levels may affect stem cell behaviour and reduce cell proliferation, contributing to the growth-restricted phenotype. High levels of stress can stimulate release of pro-inflammatory cytokines and anti-angiogenic factors, increasing the risk of pre-eclampsia. In addition, chronic stress may promote senescence of the trophoblast, which in other cell types leads to a pro-inflammatory senescence-associated secretory phenotype. Evidence from rodents suggests that a degree of trophoblastic stress develops with increasing gestational age in normal pregnancies. The increase in maternal concentrations of soluble fms-like tyrosine kinase-1 (sFlt-1) and reduction in placental growth factor (PlGF) suggest the same may occur in the human, starting around 30 weeks of pregnancy. Placental malperfusion, or co-existing maternal conditions, such as diabetes, will exacerbate that stress. Amelioration of trophoblastic stress should remain a research priority, but will be difficult due to the complexity of the molecular pathways involved.

© 2016 Elsevier Ltd. All rights reserved.

### 1. Introduction

The syncytiotrophoblast of the human placenta is a highly dynamic tissue, performing a wide range of energy-demanding functions essential for the maintenance of a successful pregnancy and normal growth of the fetus. These functions include the active transport of amino acids, ionic pumping, steroid and peptide hormone synthesis, and secretion of a wide array of growth factors and cytokines. As a consequence, the syncytiotrophoblast is equipped with a large number of mitochondria and quantities of rough endoplasmic reticulum (ER). The demands on these organelles vary depending on the stage of gestation, and in response to maternal and fetal cues. Both mitochondria and the ER must be capable of

adaptive responses, and given their roles the signalling pathways involved are central to the maintenance of cellular homeostasis. Indeed, the evidence of considerable cross-talk between the two indicates that they operate as an integrated unit. Here, we review the potential roles of these pathways in trophoblast biology, from differentiation of stem cells to senescence.

### 2. The mitochondria and endoplasmic reticulum, an integrated unit

Mitochondria are widely recognised as the major source of energy in most eukaryotic cells through their production of ATP. In the syncytiotrophoblast they have additional important functions as the site of synthesis of steroid hormones, as well as being involved in the transport and metabolism of cholesterol [1]. Mitochondria are also known to be highly dynamic organelles, forming a reticular network that undergoes continual fission and fusion, altering their morphology and with it their function [2,3]. During the

<sup>\*</sup> Corresponding author. Physiological Laboratory, Downing Street, Cambridge, CB2 3EG, UK.

E-mail address: [gjb2@cam.ac.uk](mailto:gjb2@cam.ac.uk) (G.J. Burton).

differentiation of cytotrophoblast cells prior to fusion with the syncytiotrophoblast, the mitochondria undergo morphological changes that generate smaller, more rounded profiles. Tubulovesicular cristae typical of steroidogenic cells become prevalent in the syncytiotrophoblast later in gestation [4], along with large quantities of ER. The ER is best known as the site of synthesis and post-translational processing of proteins destined for insertion into the cell membrane or for secretion. However, the ER plays additional roles as the major intracellular store of calcium, and the site of loading of peptides onto MHC class I molecules. In addition, lipogenic reactions occur on its outer surface. Perturbations of placental mitochondrial and ER homeostasis therefore have the potential to drive diverse downstream effects.

Because calcium and lipid metabolism are central to the function and regulation of both organelles, there are close physical contacts between their respective membranes for integrative signalling. Mitochondrial and ER membranes come into close apposition at punctate sites somewhat similar to synapses in the nervous system, referred to collectively as the mitochondria-associated ER membrane (MAM). Calcium transporters and ion channels are concentrated at these sites, and flux of calcium between the two organelles is thought to link their functionality in a bidirectional manner [5,6]. Thus, calcium-mediated ER signals may stimulate tricarboxylic acid (TCA) cycle activity, complexes of the mitochondrial electron transport chain and ATP production to meet the demands of protein synthesis. In turn, ER maintenance of intracellular calcium homeostasis through SERCA pumping may be responsive to cellular energy levels, and thus mitochondrial activity. Contact between the two organelles must, however, be closely controlled, since mitochondrial  $\text{Ca}^{2+}$  overload can also modulate cell death [6,7]. The ER-mitochondria interface additionally provides a platform for the regulation of a number of other shared functions (see Fig. 2 of Ref. [8]), including autophagy, inflammatory formation [8], the removal of damaged mitochondria [9], and initiation of the apoptotic machinery. A number of regulatory and chaperone proteins have been identified at MAMs, and so such functions may be responsive to changes in the microenvironment or vulnerable to perturbation by stressors [5,6]. Many of these proteins are also involved in the processes of mitochondrial fission and fusion, suggesting that modulation of mitochondrial morphology in response to stress is a further function of the ER.

In mammalian cells, optic atrophy 1 (Opa1) and the mitofusins (Mfn1 and Mfn2) bring about mitochondrial fusion, whilst dynamin 1-like (Drp1) and fission 1 (Fis1) are involved in the process of fission [10]. Mitochondrial-ER interactions are strongly implicated in playing a key role in mitochondrial fission [8], with ER tubules mediating the formation of mitochondrial constriction sites via actin polymerisation induced by the ER-associated protein inverted formin 2 (INF2) [11] (see Fig. 1 in Ref. [12]). Actin polymerisation is also proposed to enhance assembly of Drp1 complexes at the outer mitochondrial membrane, bringing about a secondary mechanism of constriction and ultimately mitochondria fission [11]. The involvement of the ER in fission might therefore explain the changes in mitochondrial morphology that accompany an increase in ER mass during the differentiation of cytotrophoblast cells, thereby enhancing capacity in energy production.

Equally, there is increasing recognition of an interaction between the mediators of mitochondrial fusion and the ER. It has been known for some time that Mfn2 mediates an interaction between the mitochondrial reticulum and the ER [13]. Recently, ablation of Mfn2 in mouse embryonic fibroblasts (MEFs) was found to result in increased connections between the ER and mitochondria, and an increased susceptibility to  $\text{Ca}^{2+}$ -overload dependent cell death [14]. It has therefore been proposed that Mfn2 acts as a tethering antagonist, preventing an excessive proximity between

the mitochondria and the rough ER [14], whilst a similar function has been proposed for Mfn1 in regulating contacts between the mitochondria and smooth ER [15]. Both Mfn1 and Mfn2 have been shown to play vital roles in embryonic development, underlining the importance of continual mitochondrial fusion. However, *Mfn2* mutant embryos also show specific and severe disruption of the placental trophoblast giant cell layer [16], which may be due to the high metabolic activity of these cells associated with polyploidy. Loss of function of this trophoblast lineage is thought to contribute to the high incidence of mid-gestation embryonic death in these mutants.

### 3. Oxidative, mitochondrial and ER stress

Attention has focussed on mitochondria within the syncytiotrophoblast, and more recently the ER, due to their involvement in placental stress. It has been recognised for some time that placental oxidative stress is associated with complications of pregnancy, such as pre-eclampsia and growth restriction [17–19], and it is thought to play a significant role in their pathophysiology.

There are many potential sources of reactive oxygen species (ROS) within the syncytiotrophoblast, such as the detoxification of drugs and xenobiotics by cytochrome P450, the response of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase to growth factors and cytokines, and various other oxido-reductase and cyclooxygenase enzymes. However, the mitochondria are considered to be a major source due to the generation of superoxide radicals through the transfer of electrons onto molecular oxygen by complexes I and III of the electron transport chain [20]. This transfer is particularly significant under hypoxic conditions [21,22]. As the superoxide anion is polarised, it remains within the mitochondrial matrix where it is enzymatically converted to hydrogen peroxide by manganese superoxide dismutase. However, excessive generation of ROS, or comprise of the defences through deficiency of catalytic micronutrients, such as selenium, can result in damage to biological molecules within the matrix. Unlike nuclear DNA, mtDNA is not protected by histones and so is vulnerable to oxidative-induced mutation. Furthermore, proteins encoded by mtDNA are synthesised within the matrix and the protein-folding environment can be disturbed by excessive ROS. Mutations and/or aberrant mitochondrial protein folding may compromise assembly of the subunits of ETC into functional complexes, thereby limiting energy production and increasing the risk of further ROS production [23]. Thus, placental oxidative stress is increased in patients with diabetes mellitus when oxidative phosphorylation is enhanced [24], and the risk of developing hypertensive disorders in pregnancy is inversely related to the maternal selenium concentration [25,26]. Mitochondria isolated from placentas of severely pre-eclamptic patients show a dramatically altered morphology [27], though the extent to which this disruption is exaggerated by the process of isolation is unclear. ROS are also generated through the shorter electron transport chain within the ER that traffics electrons during the formation of disulphide bonds [28]. Under normal physiological conditions this accounts for ~25% of ROS generation within cells, although this proportion may be higher in those with a high secretory output, such as the syncytiotrophoblast.

Because of the coupling of mitochondrial and ER function through MAMs, oxidative, mitochondrial and ER stress are closely interlinked, and each is unlikely to occur in isolation [29–31]. Thus, ROS induce calcium release from the ER through the inositol-1,4,5-triphosphate (IP3R) and ryanodine receptors, with the ions being taken up by the mitochondria at MAMs [5] stimulating further ROS production and generating a dangerous feed-forward situation. Moreover, excessive mitochondrial fission, resulting in a fragmented network, is also associated with increased ROS

production [3,32], and this may represent a protective mechanism serving to isolate and subsequently remove damaged mitochondria. Fission can result in mitochondrial depolarisation due to a loss of membrane potential ( $\Delta\psi_m$ ), targeting such fragmented mitochondria for mitophagy [33]. A number of possible mechanisms that link increased ROS production with increased fission have been proposed, although notably there have been no reports that the fission/fusion proteins are themselves directly redox-sensitive [34]. Mitochondrial fusion can act as a rescue pathway for fragmented mitochondria, including those that undergo transient depolarisation. However, if mitochondrial depolarisation is sustained (as seen if a mitochondrial uncoupler is administered) fusion is inhibited and mitophagy thus favoured [33]. It has been suggested that upregulation of pro-fusion proteins may therefore offset some of the consequences of oxidative stress [34], with a possible role for the ERK-JNK pathway implicated [35]. Such promotion of the molecular mediators of fusion (e.g. Mfn2) would also be likely to alter the interaction between mitochondria and ER, and this may in turn offer some protection under conditions of ER/oxidative stress.

Within mitochondria, ROS can also induce damage to mtDNA and may culminate in opening of the membrane permeability transition pore, leading to defective functioning of the electron transport chain, loss of  $\Delta\psi$  and a collapse in ATP production [36]. Alternatively, an adaptive antioxidant response to mitigate against increased ROS at the cost of impaired mitochondrial efficiency may occur via the upregulation of the mitochondrial uncoupling protein, UCP2. UCP2 expression increases towards term in the human placenta [37], and does so similarly in the rat placenta in association with the expression of other antioxidant enzymes [38]. Mild uncoupling leading to a slight loss of  $\Delta\psi$  is believed to be protective against ROS, with UCP2 implicated in cytoprotection in a number of tissues, but controversy still surrounds the question of whether UCP2 exhibits uncoupling activity under physiological conditions, not least because this is technically difficult to establish [39]. Moreover, there is presently no evidence to support an antioxidant role for UCP2 in the human placenta [37].

Defective mitochondrial function may promote accumulation of unfolded and misfolded proteins within the matrix, initiating stress. Similarly, loss of calcium homeostasis within the ER impairs the protein folding machinery, resulting in the accumulation of unfolded or misfolded proteins in the lumen, a condition that constitutes ER stress.

#### 4. The unfolded protein response

Protein synthesis is core to many cellular events, and so its regulation is integral to homeostatic mechanisms [40]. In eukaryotic cells, protein synthesis occurs in the cytosol, endoplasmic reticulum and mitochondria. In each compartment, a conserved group of organelle-specific molecular chaperones facilitate efficient folding of nascent, unfolded polypeptides into their final, distinctive conformation. Stress occurs when accumulated unfolded or misfolded proteins exceed the compartment's folding capacity, posing the risk of catastrophic damage. To cope with this risk, organelle-specific signalling pathways, known as unfolded protein responses (UPRs), have evolved. These pathways aim to restore homeostasis by promoting expression of compartment-specific genes to increase molecular chaperones and folding capacity, as well as genes involved in protein degradation to remove the accumulated toxic misfolded proteins.

The protein-folding environment of the ER is guarded by the UPR<sup>ER</sup>, which comprises three signal transduction pathways [41]. They are the PERK (PKR-like endoplasmic reticulum kinase), ATF6 (activating transcription factor 6) and IRE-1 (inositol-requiring enzyme-1) pathways (Fig. 1). The sensors are located on the luminal

side of the ER membrane and are activated by the accumulation of misfolded proteins. PERK is a kinase with two principal targets. Firstly, it phosphorylates eukaryotic initiation factor 2- $\alpha$  (eIF2- $\alpha$ ), providing a rapid block to translation and reducing the burden on the folding machinery. Secondly, it activates the transcription factors NRF2 and ATF4; the former directly and the latter through phosphorylation of eIF2- $\alpha$ . The ATF6 and IRE-1 pathways also lead to activation of transcription factors, ATF6 and XBP-1 respectively (Fig. 1). Together, these four transcription factors co-ordinate expression of an array of genes that enhance antioxidant defences and increase the folding capacity of the ER. Thus, uptake of cystine and its conversion to glutathione is stimulated, along with expression of ER chaperone proteins, haemoxygenase enzymes, lipogenesis and synthesis of more ER cisternae [42–44]. If these responses fail to restore ER and redox homeostasis, activation of apoptotic pathways occurs through expression of the protein CHOP, and direct signalling through caspase 4.

Trophoblast cell lines show a graded response to exogenous inducers of ER stress, such as tunicamycin that blocks *N*-glycosylation of proteins or thapsigargin that disrupts calcium homeostasis. Low doses cause only increased phosphorylation of eIF2- $\alpha$ , while higher doses increase the chaperone proteins and finally activate the CHOP pathway [45]. Teleologically, apoptosis should only be induced as an action of last resort if homeostasis cannot be restored.

Although the UPR pathways evolved to regulate ER function, they have subsequently been recruited to modulate other cellular activities [40]. ChIP-seq analysis in skeletal muscle has revealed that approximately 40% of XBP-1 targets are unrelated to ER function, including, for example, genes associated with myogenic differentiation [46]. Hence, ER stress may have diverse effects.

Increasing evidence suggests the existence of UPR pathways in mitochondria too, UPR<sup>mt</sup>, which are particularly active in cells with a high rate of biogenesis, high ROS production, and defective mitochondrial structure [47–49]. Conceptually similar to the UPR<sup>ER</sup>, once mitochondria sense unfolded or misfolded proteins, a signal is transmitted to the nucleus, promoting expression of genes encoding mitochondria-specific molecular chaperones and quality-control proteases in order to re-establish mitochondrial homeostasis [50]. The signalling pathways involved in the UPR<sup>mt</sup> are not well understood. It has been demonstrated that a quality control protease, ClpP, which is located within the mitochondrial matrix, and the ATP Binding Cassette (ABC) transporter, which resides in the mitochondrial inner membrane, may play key roles. ClpP recognizes and degrades misfolded proteins into short peptides, which in turn are extruded into the intermembrane space by the ABC transporter. Knock-down of the *ClpP* gene prevents mitochondrial stress-induced expression of mitochondrial molecular chaperone genes [50]. Equally, deletion of the ABC transporter orthologous gene, *HAF-1* in *C. elegans*, attenuates UPR<sup>mt</sup> activation upon mitochondrial stress, suggesting ClpP-derived peptides may act as signalling components in the UPR<sup>mt</sup> [51]. The UPR<sup>mt</sup> activates the bZip transcription factor ATFS-1 (Activating Transcription Factor associated with Stress, also known as ZC376.7), which translocates to the nucleus to stimulate expression of mitochondrial molecular chaperone genes, including mitochondrial heat shock protein 70 (mtHSP70) and HSP60. The transcriptional regulatory function of ATFS-1 requires interaction with another transcription factor, homeobox transcription factor DVE-1, and with co-factor ubiquitin-like protein UBL-5 [50].

#### 5. Involvement of UPR pathways in placental physiology and development

Activation of the UPR pathways should not necessarily be




Fig. 1. Diagrammatic representation of the three signalling arms of the UPR<sup>ER</sup> and their principal downstream effectors.

interpreted as evidence of pathology, and may be a cell-type and condition-specific response to normal events, such as differentiation or variations in physiology. Thus, the pancreas and placenta display low-grade activation of the UPR<sup>ER</sup> under normal conditions because of their high endocrine and exocrine activity [52,53]. Another example is in the crypts of the intestine, where markers of activation of the UPR<sup>ER</sup> pathways only become apparent as the stem cells differentiate into transit amplifying cells, presumably reflecting a change in the requirements for cell surface and secreted proteins. Administration of thapsigargin to organoid cultures of these stem cells causes loss of stemness in a PERK-eIF2- $\alpha$  dependent manner, indicating the importance of UPR<sup>ER</sup> pathways for stem cell behaviour [54]. Finally, recent evidence indicates mitochondrial stress signaling induces cellular adaptations that reduce the impact of subsequent exposure to lethal stressors [55].

Although there is considerable overlap in the downstream targets of the three arms of the UPR<sup>ER</sup>, they also have unique functions. Thus, it might be expected that in some circumstances components of the UPR<sup>ER</sup> are activated selectively. This is seen during the differentiation of plasma cells when the synthesis of antibodies is upregulated. There is clearly a need to increase the capacity of the ER as part of this process, but it would be counterproductive to inhibit protein translation at the same time. Hence, plasma cell differentiation is associated with stimulation of the IRE-1 and ATF6 pathways, but no phosphorylation of eIF2- $\alpha$  [56].

Limited data are available for the role of these pathways during development of the placenta. Using a transgenic reporter mouse, Iwawaki et al. demonstrated activation of the IRE-1 pathway during normal development at E14.5 [53]. By contrast, activation in the embryo was minimal. Genetic knock-out of the pathway causes aberrant development of the placenta, particularly in the labyrinth zone where there is abnormal vascularisation associated with reduced levels of VEGF. This is accompanied by reduced proliferation of trophoblast cells, but no increase in apoptosis [53]. Overall, it is apparent that placental changes are responsible for the increase in embryonic deaths observed in the mutant mice.

These findings indicate that UPR signalling pathways are important during placental development. The converse is also true, that elevation of ER and mitochondrial stress can perturb normal formation of the organ. In a mouse model of constitutive ER stress due to a dysfunctional mutation in eIF2- $\alpha$ , there is evidence of premature differentiation of trophoblast stem cells at E9.5 [57]. Embryonic fibroblasts isolated from the mutants proliferate at a rate 50% slower than in controls. Activation of the PERK pathway and accumulation of aberrantly glycosylated proteins are only observed in the endocrine junctional zone [57]. Hence, perturbation of ER function in the murine placenta is context-dependent, and predominantly affects cells with a high secretory output. It is notable that pups homozygous for the mutation die post-natally due to severe defects in pancreatic function [58], suggesting there are parallels between the junctional zone and the pancreas.

Administration of tunicamycin to pregnant dams has also been used to test the effects of prolonged ER stress on placental development [59]. This results in reduced placental weight and disruption of placental morphogenesis, with abnormal interdigitation between the labyrinth and junctional zones. There is also reduced vascularisation of the labyrinth zone, and reduced expression of *Slc2a1*, the GLUT1 glucose transporter, but increased expression of *Slc2a3*, the GLUT3 transporter.

There are few data available at present that implicate the UPR<sup>mt</sup> in placental development. The UPR<sup>mt</sup> can affect cell cycle proteins, slowing cell proliferation or even inhibiting replication of cells. Cells containing defective mitochondrial structures have altered expression of many cell cycle regulators including p19, a cyclin-dependent kinase inhibitor involved in cell cycle arrest at G1 [60]. There is a reduction in the expected number of *Clpp*-deficient progenies from heterozygous crosses, indicating increased embryonic loss [61]. It is unclear whether fetal growth restriction occurs in these animals, although *Clpp*-deficient offspring exhibit smaller body size in adulthood [61].

Hence, experimental evidence suggests that UPR pathways play physiological roles in placental development. Oxidative stress has



also been implicated in placental development, but more in terms of stimulating villous regression than proliferation. Towards the end of the first trimester the villi that initially form over the entire surface of the human chorionic sac regress over the superficial pole, leaving the definitive discoid placenta. Regression is temporally and spatially associated with onset of the maternal circulation, and locally high levels of oxidative stress are thought to suppress villous proliferation and induce apoptosis [62,63]. As this process occurs in all pregnancies it is considered physiological.

## 6. Involvement of UPR pathways in placental adaptive responses

The boundary between physiological and pathological events is inevitably blurred, with these labels being at opposite ends of a spectrum. A pertinent example is the response to chronic hypobaric hypoxia experienced during pregnancy at high altitude. Placentas from normal pregnancies in women of non-indigenous descent at 3100 m in Colorado show increased phosphorylation of eIF2- $\alpha$  compared to sea-level controls, but no other evidence of activation of the UPR<sup>ER</sup> [64]. Additionally, many subunits of ETC complexes were found to be down-regulated without a change in mitochondrial density, indicating defective mitochondrial function and potential activation of UPR<sup>mt</sup> [65]. These findings can be replicated by exposing placental cells to hypoxia in culture, and is associated with reduced cell proliferation. Treatment of normoxic cells with salubrinal, a p-eIF2- $\alpha$  phosphatase inhibitor, also suppresses proliferation, and reduces expression of subunits in ETC complexes, confirming that phosphorylation of eIF2- $\alpha$  alone is sufficient to account for the reduced volume of the placental villous trees measured in these placentas [64,66]. A similar increase in phosphorylation of eIF2- $\alpha$  and reduction in mitochondrial ETC subunits occurs in the murine placenta when dams are maintained under 13% oxygen throughout gestation, equivalent to pregnancy at 3100 m [67]. In both cases there is mild growth restriction of the fetus, and so the changes may be viewed as an adaptive response that matches growth to oxygen availability.

Attenuation of protein translation is a part of a cell's repertoire of adaptive responses to cope with hypoxia [68], for incorporation of a single amino acid into a nascent polypeptide chain requires four high-energy phosphate bonds. The consequences will depend on the duration of the attenuation and the half-life of the proteins concerned, but we have observed further links between ER stress and mitochondrial function through this mechanism. Thus complexes of the mitochondrial electron transport chain are reduced at the protein, but not mRNA, level in the high-altitude placenta, consistent with the reduction in ATP/ADP ratio observed [65,69]. This effect may have been mediated through the PERK-eIF2- $\alpha$  pathway, for administration of salubrinal to trophoblast-like cells is sufficient to reduce the complexes at the protein level, with a particular stark loss of complex I [65]. While reducing mitochondrial activity might at first sight be considered an adverse response, it is potentially beneficial as long as it is matched by reduced energy demands through translational arrest and/or an increased reliance on glycolysis for ATP production [70–72]. Production of ROS at complexes I and III is paradoxically increased during hypoxia due to an accumulation of electrons on the electron transport chain in the absence of oxygen as the final electron acceptor [21,22,70]. By reducing these complexes, cells may protect themselves from excessive generation of ROS under these conditions. It is notable that a similar reduction in electron transport chain complexes is observed in skeletal muscle of climbers at high altitude [73,74] in association with an altered energetic profile [75].

## 7. Involvement of UPR pathways in placental pathology

Many common, non-infective complications of pregnancy are associated with deficient physiological conversion of the maternal spiral arteries supplying the placenta [76]. It is frequently asserted that this leads to placental hypoxia, although no measurements have been recorded from the intervillous space *in vivo* to confirm or refute this supposition. Whether the problem is one of hypoxia or ischaemia-reperfusion-type injury has yet to be resolved [19], but there is conclusive evidence of oxidative stress in placentas from patients suffering fetal growth restriction and pre-eclampsia. Activation of the UPR<sup>ER</sup> is seen in growth-restricted placentas of maternal vascular origin, and in placentas from cases of early-onset, but not late-onset, pre-eclampsia [45,77]. The level of activation is greater, and involves more pathways, than is seen in the normal placentas from high altitude, indicating a more severe insult. In the growth-restricted placentas, levels of AKT were reduced at the protein, but not mRNA, level due to translational arrest, along with reduced cyclin D1. When comparing with gestationally age-matched placentas from pregnancies where the growth restriction was complicated by pre-eclampsia, the levels p-eIF2- $\alpha$  and ER chaperone proteins were even higher, and there was additional activation of the CHOP pro-apoptotic pathway [45].

In comparison, the evidence of UPR<sup>mt</sup> involvement in pathological pregnancies is very limited. There is an increase of mitochondrial molecular chaperone HSP60 immunoreactivity in the syncytiotrophoblast and cytotrophoblast cells of growth-restricted placentas, especially in regions of thrombi, syncytial knots and avascular villi [78], but no other data.

The changes in the AKT-mTOR pathway, a central regulator of cell proliferation, and in cyclin D1 are sufficient to explain the growth restricted phenotype, but can differences in the placental pathophysiology explain the superimposition of pre-eclampsia on this phenotype? Pre-eclampsia, especially the early-onset form, is an inflammatory state in which elevated serum concentrations of anti-angiogenic and pro-inflammatory cytokines combine to cause maternal endothelial activation. It is generally assumed that the placenta is the source of these factors, as delivery is followed by a rapid decline in their levels, although a contribution from maternal sources cannot be ruled out. Generation of oxidative stress in the placenta, either *in vitro* through hypoxia-reoxygenation or *in vivo* through the ischaemia-reperfusion that accompanies labour, mimics the transcriptional changes seen in pre-eclampsia. It also stimulates secretion of pro-inflammatory cytokines and anti-angiogenic factors, including soluble fms-like tyrosine kinase-1, sFlt-1, which has been implicated in the pathophysiology of pre-eclampsia [79–82]. Secretion is mediated through the p38, NF $\kappa$ -B, and stress-activated protein kinase MAPK pathways [83].

Again, there are multiple links between the UPR<sup>ER</sup> and these pathways (Fig. 2). Although IRE-1 is generally considered an endoribonuclease that splices *XBP-1* mRNA to yield a functional transcription factor, at high levels of activation it has an additional kinase activity and is capable of phosphorylating TRAF2. This in turn activates the p38, NF $\kappa$ -B pathways and JNK [84,85]. Furthermore, attenuation of protein translation leads to activation of NF $\kappa$ B, as the half-life of the inhibitory sub-unit, I $\kappa$ B, is shorter than that of NF $\kappa$ B [86]. Hence, it is not unreasonable to envisage that the obstetric outcome is dependent on the severity of the UPR<sup>ER</sup> response, which in turn may reflect interactions between the degree of the vascular insult and maternal factors, such as her genotypic predisposition to endothelial disease and the state of her antioxidant defences. In support of this hypothesis, the few data available indicate that spiral arterial conversion is most deficient in cases of growth restriction associated with pre-eclampsia [87]. Magnetic resonance imaging has also demonstrated reduced perfusion of the




Fig. 2. Summary of the mitochondrial and ER pathways that may contribute to the activation of proinflammatory pathways in the syncytiotrophoblast under conditions of stress.

placenta in cases of early-onset but not late-onset pre-eclampsia [88], and a change in the phosphodiester/phosphomonoester ratio of the  $^{31}\text{P}$  signal indicates accelerated ageing of the tissues [89].

### 8. Syncytiotrophoblast senescence

One of the consequences of chronic oxidative, mitochondrial and ER stress is that they can induce senescent changes in tissues [60,90,91]. Senescence is characterised by irreversible cell-cycle arrest, cytological and metabolic changes and the acquisition of a senescent-associated secretory phenotype (SASP) that leads to the release of a mix of pro-inflammatory cytokines and proteases. Whilst cause and effect can be difficult to establish, senescence is strongly associated with a number of mitochondrial perturbations, which in turn share a connection with ER stress. These include excessive ROS production, increased mitochondrial fusion, mitochondrial uncoupling and depolarisation of the inner mitochondrial membrane, loss of ATP and activation of AMPK, decreased  $\text{NAD}^+$  availability and mitochondrial  $\text{Ca}^{2+}$  accumulation [10], while pharmacological inhibition of electron transport chain complexes I, II or III can also lead to premature senescence [10,92–94].

Senescence has only recently been considered as a biological phenomenon in the syncytiotrophoblast [95,96], although changes associated with ageing have long been recognised morphologically [97,98]. It has been suggested that the retrovirally-driven fusion process by which differentiated cytotrophoblast cells are incorporated into the syncytiotrophoblast initiates the process, and molecular markers of senescence are displayed by the syncytiotrophoblast of mature, but otherwise normal, placentas [95]. Certainly, mitosis has never been reported in the syncytiotrophoblast, and aggregations of nuclei displaying condensed chromatin and evidence of oxidative damage accumulate as gestation advances [99]. Therefore, it may be the case that the syncytiotrophoblast undergoes a degree of molecular senescence in all pregnancies, triggered perhaps by increasing stress arising from

a progressive mismatch between maternal perfusion and fetal demands [100] (Fig. 3). This is difficult to prove as longitudinal sampling of the human placenta cannot be performed, but circumstantial evidence from circulating biomarkers indicative of placental well-being suggest that this may be the case. For example, maternal concentrations of cell-free fetal DNA, which can be released by the placenta when subjected to oxidative stress [101], increase steeply after 30 weeks of gestation [102].




Fig. 3. Schematic representation of changes in oxygen concentration in the intervillous space (IVS) of the placenta, fetal weight, and maternal concentrations of sFlt-1 and PIGF across gestational age. A burst of oxidative stress is observed in the trophoblast at the end of the first trimester, associated with onset of the maternal arterial circulation and remodelling of the early placenta into the definitive form. Placental secretion of sFlt-1 is positively regulated by oxidative stress, while that of PIGF is negatively regulated by ER stress. The changes in maternal concentrations may reflect an increase in stress within the trophoblast towards term induced by a progressive mismatch in fetal demand for oxygen and maternal supply. This may be exacerbated in cases of early-onset pre-eclampsia (dashed line) due to malperfusion secondary to deficient remodelling of the spiral arteries. Chronic stress may induce senescent changes in the trophoblast, but the point at which that occurs is uncertain.

Longitudinal studies in rodent models have yielded more solid evidence. An increase in oxidative stress is observed during late gestation in the labyrinth zone that performs gaseous exchange in the mouse [103], and is associated with an increase in the mitochondrial DNA copy number [104]. Similar changes in oxidative stress have been reported in the rat [38].

Increasing stress within the syncytiotrophoblast during the third trimester could explain the gradual increase in maternal serum concentration of sFlt-1 from 29 to 32 weeks onwards [105]. In addition, maternal concentrations of placental growth factor (PlGF) decline after the same time-point. This factor is negatively regulated in trophoblast-like cell lines by the ATF4 and ATF6 pathways [106], and so again the data are consistent with increasing placental stress. Thus, it may be that the changes that occur in complications of pregnancy are an exaggeration of normal placental ageing, induced by malperfusion secondary to deficient remodelling of the spiral arteries (Fig. 3). In support of this hypothesis, shortening of telomeres, a hallmark of senescence, is greater in placentas from pregnancies complicated by growth restriction and pre-eclampsia than in normal controls [107–109].

Many components of the SASP, including IL-1, IL-6, IL-8, are increased in early-onset pre-eclampsia and contribute to the maternal inflammatory state. The inflammatory component of the SASP is regulated principally through High-mobility group box 1 (HMGB1) and the NF $\kappa$ B pathway [110], demonstrating the overlap among the oxidative stress, UPR and senescence signalling cascades. HMGB1 signalling is increased in the syncytiotrophoblast in cases of severe pre-eclampsia [111], creating what might be considered a sterile inflammatory response. In normal situations, the function of the SASP is to attract immune cells that will remove the senescent one. In the case of the syncytiotrophoblast that is not possible due to its unique syncytial nature. Consequently, the response may have been preserved in the cellular machinery, but has the detrimental effects of activating maternal immune and endothelial cells, resulting in the syndrome of pre-eclampsia. Viewing the pathophysiology in this way may open new avenues for therapeutic interventions.

## 9. Autophagy in the placenta

One of the protective mechanisms cells can deploy against senescence is autophagy, whereby aggregates of misfolded proteins or damaged organelles are degraded through the lysosomal pathway and their constituent elements recycled. There is increasing evidence that prolonged activation of the UPR<sup>mt</sup> ultimately initiates mitophagy of damaged mitochondria [112,113]. Additionally, there are also close links between the UPR<sup>ER</sup> pathways, in particular the PERK and IRE arms, and various components of the autophagocytic machinery [30,114,115]. It might be expected, therefore, that autophagy occurs in the syncytiotrophoblast in pre-eclamptic or growth restricted pregnancies, but the evidence from different studies is conflicting. Most have relied on changes in the levels of key regulators, such as LC3-II, LAMP-2 and beclin-1, at the mRNA or protein level in placental homogenates as markers. On this basis, increased autophagy has been reported in placentas from pregnancies complicated by growth restriction, with or without pre-eclampsia, but not from those with pre-eclampsia alone [116,117]. Conversely, other studies have found evidence of an increase in placentas from pregnancies complicated by hypertensive disorders, independent of the presence of growth restriction [118,119]. Few studies have included electron micrographs localising the phenomenon, but those available indicate that it occurs in the syncytiotrophoblast and to a lesser extent in cytotrophoblast cells, and involves mitochondria in growth-restricted pregnancies [116–118]. Whilst autophagy can be induced in primary cultures of

cytotrophoblast or trophoblast cell lines *in vitro* through exposure to hypoxia and/or glucose deprivation [120–122], the significance of this response for placental function and fetal well-being *in vivo* remains uncertain. Nonetheless, the possibility remains that the placenta may act as a nutritional reserve that can be mobilised to protect growth of the fetus, in particular the brain, under conditions of acute deprivation [123].

## 10. Apoptosis of trophoblast

If homeostatic pathways fail to restore metabolic equilibrium, then oxidative, mitochondrial and ER stress can induce apoptosis in many cell types through activation of both mitochondrial-dependent and -independent apoptotic machineries, including the caspase pathway and CHOP. For the syncytiotrophoblast this presents a major threat, as in the absence of cell boundaries there is a danger of apoptosis sweeping through the whole epithelium, leading to pregnancy failure. Such a wave has been observed in *in vitro* models of the syncytium [124], but *in vivo* the syncytiotrophoblast appears resistant to apoptosis, possibly to prevent such a catastrophic event [125]. By contrast, cytotrophoblast cells are vulnerable to both apoptotic and necrotic cell death [125,126], and the incidence is increased in complications of pregnancy, such as miscarriage, pre-eclampsia and growth restriction [127,128], when it may reflect elevated levels of stress. Excessive loss of cytotrophoblast cells through this mechanism may have detrimental consequences for placental function by compromising the capacity for regeneration of the syncytiotrophoblast through fusion. Recruitment of cytotrophoblast cells into the syncytium appears to continue through to term [129], and as well as expanding the tissue it brings in fresh mitochondria, ER, and other organelles that may replace aged or damaged examples removed through autophagy. Unfortunately, there are no data available to indicate the rate of such turnover in the syncytiotrophoblast of either healthy or pathological placentas *in vivo*.

## 11. Broader implications of ER and oxidative stress for placental pathology

Most attention with regards to oxidative and ER stress has focused on the placenta and the villous trophoblast, but there are reports of increased stress in the decidua in pathological pregnancies. Activation of the PERK-eIF2- $\alpha$  and ATF6 pathways has been reported in decidual cells, extravillous trophoblast and macrophages in cases of early-onset growth restriction with and without pre-eclampsia, but not in cases of pre-eclampsia alone [130]. Increased oxidative and ER stress has also been observed in the decidua of cases of early pregnancy loss [131]. In both situations it is difficult to distinguish between cause and effect, but stress responses may impair trophoblast invasion or induce excessive apoptosis, compromising spiral artery remodelling.

In addition, it is possible that ER stress may affect interactions with the maternal immune cells. Extravillous trophoblast cells express the non-polymorphic class I antigens HLA-G and HLA-E, and also the highly polymorphic HLA-C. Interactions between HLA-C and the killer immunoglobulin-like receptors on the uterine natural killer cells are crucial for a successful pregnancy, and indeed appear to regulate birth weight across the natural range [132]. In particular, it is necessary to have a sufficient degree of activation of the natural killer cells, which is thought to mediate their release of proteases and cytokines necessary for remodelling of the maternal arteries. Since peptides are loaded on to MHC molecules within the ER lumen it is possible that loss of ER homeostasis in an extravillous trophoblast cell may compromise its antigenic profile. It is notable that treatment of thyroid cells with thapsigargin or tunicamycin

reduced MHC class I expression and was associated with increased natural killer cell cytotoxicity [133]. One might speculate, therefore, that induction of ER stress in the invading extravillous trophoblast, possibly induced by low-grade inflammation within the decidua or maternal metabolic disorders, might compromise activation of the uterine natural killer cells, so impairing maternal arterial remodelling with the end result of placental malperfusion and growth restriction and/or pre-eclampsia.

## 12. Conclusion

Mitochondria and the endoplasmic reticulum are two of the most dynamic and important cell organelles, performing functions central to homeostasis, viability, and growth. Their functional interdependence requires that their activities are closely interlinked, which is achieved through bidirectional signalling at MAMs. This signalling is so extensive they may be considered physiologically as single unit, the function of which may be perturbed by changes in oxygenation, glucose availability and other environmental cues. The extensive complement of both organelles in the syncytiotrophoblast necessary to meet its high metabolic and synthetic activities means that the tissue is vulnerable to oxidative and ER stress. Evidence from rodent species indicates that trophoblastic stress increases with gestational age, and so a degree of stress may be a feature of all otherwise healthy, mature placentas. High levels of trophoblastic stress are associated with complications of pregnancy, and attenuation of protein synthesis and aberrant secretion of pro-inflammatory cytokines and anti-angiogenic factors may contribute to the pathophysiology of growth restriction and pre-eclampsia respectively. Chronic stress may also promote trophoblast senescence, which in turn leads to the secretion pro-inflammatory factors that may further contribute to the pre-eclamptic syndrome. Attempts to reduce trophoblastic stress should therefore remain a research priority, but the complexity of the interactions between the mitochondria and endoplasmic reticulum require that a holistic approach to restore homeostasis be adopted rather than targeting any particular individual pathway.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## Acknowledgements

The authors thank all their colleagues and collaborators who have contributed to their studies cited here. GJB gratefully acknowledges the support of the Wellcome Trust (069027/Z/02/Z and 084804/2/08/Z), the MRC (G9701485) and Tommy's (Grant 41); AJM thanks the Research Councils UK, Action Medical Research (SP4545) and the WYNG Foundation of Hong Kong.

## References

- [1] F. Martinez, S. Olvera-Sanchez, M. Esparza-Perusquia, E. Gomez-Chang, O. Flores-Herrera, Multiple functions of syncytiotrophoblast mitochondria, *Steroids* 103 (2015) 11–22.
- [2] D.C. Chan, Mitochondria: dynamic organelles in disease, aging, and development, *Cell* 125 (7) (2006) 1241–1252.
- [3] M. Picard, T. Taivassalo, G. Gousspillou, R.T. Hepple, Mitochondria: isolation, structure and function, *J. Physiol.* 589 (Pt 18) (2011) 4413–4421.
- [4] C.J.P. Jones, H. Fox, Ultrastructure of the normal human placenta, *Electron Microsc. Rev.* 4 (1991) 129–178.
- [5] T. Hayashi, R. Rizzuto, G. Hajnoczky, T.P. Su, MAM: more than just a housekeeper, *Trends Cell Biol.* 19 (2) (2009) 81–88.
- [6] R. Bravo, T. Gutierrez, F. Paredes, D. Gatica, A.E. Rodriguez, Z. Pedrozo, et al., Endoplasmic reticulum: ER stress regulates mitochondrial bioenergetics, *Int. J. Biochem. Cell Biol.* 44 (1) (2012) 16–20.
- [7] G. Szabadkai, M.R. Duchon, Mitochondria: the hub of cellular Ca<sup>2+</sup> signaling, *Physiology* 23 (2008) 84–94.
- [8] S. Marchi, S. Patergnani, P. Pinton, The endoplasmic reticulum-mitochondria connection: one touch, multiple functions, *Biochim. Biophys. Acta* 1837 (4) (2014) 461–469.
- [9] B. Westermann, Mitochondrial dynamics in model organisms: what yeasts, worms and flies have taught us about fusion and fission of mitochondria, *Semin. Cell Dev. Biol.* 21 (6) (2010) 542–549.
- [10] D.V. Ziegler, C.D. Wiley, M.C. Velarde, Mitochondrial effectors of cellular senescence: beyond the free radical theory of aging, *Aging Cell.* 14 (1) (2015) 1–7.
- [11] F. Korobova, V. Ramabhadran, H.N. Higgs, An actin-dependent step in mitochondrial fission mediated by the ER-associated formin INF2, *Science* 339 (6118) (2013) 464–467.
- [12] A. Kasahara, L. Scorrano, Mitochondria: from cell death executioners to regulators of cell differentiation, *Trends Cell Biol.* 24 (12) (2014) 761–770.
- [13] O.M. de Brito, L. Scorrano, Mitofusin 2 tethers endoplasmic reticulum to mitochondria, *Nature* 456 (7222) (2008) 605–610.
- [14] R. Filadi, E. Greotti, G. Turacchio, A. Luini, T. Pozzan, P. Pizzo, Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria coupling, *Proc. Natl. Acad. Sci. U. S. A.* 112 (17) (2015) E2174–E2181.
- [15] P.T. Wang, P.O. Garcin, M. Fu, M. Masoudi, P. St-Pierre, N. Pante, et al., Distinct mechanisms controlling rough and smooth endoplasmic reticulum contacts with mitochondria, *J. Cell Sci.* 128 (15) (2015) 2759–2765.
- [16] H. Chen, S.A. Detmer, A.J. Ewald, E.E. Griffin, S.E. Fraser, D.C. Chan, Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic development, *J. Cell Biol.* 160 (2) (2003) 189–200.
- [17] C.A. Hubel, Oxidative stress in the pathogenesis of preeclampsia, *Proc. Soc. Exp. Biol. Med.* 222 (3) (1999) 222–235.
- [18] L. Myatt, Review: Reactive oxygen and nitrogen species and functional adaptation of the placenta, *Placenta* 31 (Suppl:S66–9) (2010).
- [19] G.J. Burton, E. Jauniaux, Oxidative stress, *Best. Pract. Res. Clin. Obstet. Gynaecol.* 25 (3) (2011) 287–299.
- [20] E. Cadenas, K.J.A. Davies, Mitochondrial free radical generation, oxidative stress, and aging, *Free Radic. Biol. Med.* 29 (2000) 222–230.
- [21] R.D. Guzy, B. Hoyos, E. Robin, H. Chen, L. Liu, K.D. Mansfield, et al., Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing, *Cell Metab.* 1 (6) (2005) 401–408.
- [22] R.D. Guzy, P.T. Schumacker, Oxygen sensing by mitochondria at complex III: the paradox of increased reactive oxygen species during hypoxia, *Exp. Physiol.* 91 (5) (2006) 807–819.
- [23] N.G. Larsson, Somatic mitochondrial DNA mutations in mammalian aging, *Annu. Rev. Biochem.* 79 (2010) 683–706.
- [24] F. Lyall, J.L. Gibson, I.A. Greer, D.E. Brockman, A.L. Eis, L. Myatt, Increased nitrotyrosine in the diabetic placenta: evidence for oxidative stress, *Diabetes Care* 21 (10) (1998) 1753–1758.
- [25] M.P. Rayman, S.C. Bath, J. Westaway, P. Williams, J. Mao, J.J. Vanderlelie, et al., Selenium status in UK pregnant women and its relationship with hypertensive conditions of pregnancy, *Br. J. Nutr.* (2015) 1–10.
- [26] M. Xu, D. Guo, H. Gu, L. Zhang, S. Lv, Selenium and preeclampsia: a systematic review and meta-analysis, *Biol. Trace Elem. Res.* (2015) [Epub ahead of print].
- [27] Z. Shi, W. Long, C. Zhao, X. Guo, R. Shen, H. Ding, Comparative proteomics analysis suggests that placental mitochondria are involved in the development of pre-eclampsia, *PLoS One* 8 (5) (2013) e64351.
- [28] B.P. Tu, J.S. Weissman, Oxidative protein folding in eukaryotes: mechanisms and consequences, *J. Cell Biol.* 164 (3) (2004) 341–346.
- [29] K. Zhang, Integration of ER stress, oxidative stress and the inflammatory response in health and disease, *Int. J. Clin. Exp. Med.* 3 (1) (2010) 33–40.
- [30] D. Senft, Z.A. Ronai, UPR, autophagy, and mitochondria crosstalk underlies the ER stress response, *Trends Biochem. Sci.* 40 (3) (2015) 141–148.
- [31] L. Galluzzi, O. Kepp, C. Trojel-Hansen, G. Kroemer, Mitochondrial control of cellular life, stress, and death, *Circ. Res.* 111 (9) (2012) 1198–1207.
- [32] T. Yu, J.L. Robotham, Y. Yoon, Increased production of reactive oxygen species in hyperglycemic conditions requires dynamic change of mitochondrial morphology, *Proc. Natl. Acad. Sci. U. S. A.* 103 (8) (2006) 2653–2658.
- [33] G. Twig, O.S. Shirihai, The interplay between mitochondrial dynamics and mitophagy, *Antioxid. Redox. Signal* 14 (10) (2011) 1939–1951.
- [34] P. Jezek, L. Plecita-Hlavata, Mitochondrial reticulum network dynamics in relation to oxidative stress, redox regulation, and hypoxia, *Int. J. Biochem. Cell Biol.* 41 (10) (2009) 1790–1804.
- [35] O.M. de Brito, L. Scorrano, Mitofusin 2: a mitochondria-shaping protein with signaling roles beyond fusion, *Antioxid. Redox. Signal* 10 (3) (2008) 621–633.
- [36] A.J. Kowaltowski, R.F. Castilho, A.E. Vercesi, Mitochondrial permeability transition and oxidative stress, *FEBS Lett.* 495 (1–2) (2001) 12–15.
- [37] M.J. Stark, N.A. Hodyl, M. Butler, V.L. Clifton, Localisation and characterisation of uncoupling protein-2 (UCP2) in the human preterm placenta, *Placenta* 33 (12) (2012) 1020–1025.
- [38] M.L. Jones, P.J. Mark, J.L. Lewis, T.A. Mori, J.A. Keelan, B.J. Waddell, Antioxidant defenses in the rat placenta in late gestation: increased labyrinthine expression of superoxide dismutases, glutathione peroxidase 3, and uncoupling protein 2, *Biol. Reprod.* 83 (2) (2010) 250–260.
- [39] M. Donadelli, I. Dando, C. Fiorini, M. Palmieri, UCP2, a mitochondrial protein regulated at multiple levels. Cellular and molecular life sciences, *CMLS* 71 (7)



- (2014) 1171–1190.
- [40] D.T. Rutkowski, R.S. Hegde, Regulation of basal cellular physiology by the homeostatic unfolded protein response, *J. Cell Biol.* 189 (5) (2010) 783–794.
- [41] D. Ron, P. Walter, Signal integration in the endoplasmic reticulum unfolded protein response, *Nat. Rev. Mol. Cell Biol.* 8 (7) (2007) 519–529.
- [42] S.B. Cullinan, J.A. Diehl, Coordination of ER and oxidative stress signaling: the PERK/Nrf2 signaling pathway, *Int. J. Biochem. Cell Biol.* 38 (3) (2006) 317–332.
- [43] J. Lewerenz, P. Maher, Basal levels of eIF2 $\alpha$  phosphorylation determine cellular antioxidant status by regulating ATF4 and xCT expression, *J. Biol. Chem.* 284 (2) (2009) 1106–1115.
- [44] J.G. Dickhout, R.E. Carlisle, D.E. Jerome, Z. Mohammed-Ali, H. Jiang, G. Yang, et al., Integrated stress response modulates cellular redox state via induction of cystathionine gamma-lyase: cross-talk between integrated stress response and thiol metabolism, *J. Biol. Chem.* 287 (10) (2012) 7603–7614.
- [45] H.W. Yung, S. Calabrese, D. Hynx, B.A. Hemmings, I. Cetin, D.S. Charnock-Jones, et al., Evidence of placental translation inhibition and endoplasmic reticulum stress in the etiology of human intrauterine growth restriction, *Am. J. Pathol.* 173 (2008) 451–462.
- [46] D. Acosta-Alvarez, Y. Zhou, A. Blais, M. Tsikitis, N.H. Lents, C. Arias, et al., XBP1 controls diverse cell type- and condition-specific transcriptional regulatory networks, *Mol. Cell.* 27 (1) (2007) 53–66.
- [47] R.D. Martinus, G.P. Garth, T.L. Webster, P. Cartwright, D.J. Naylor, P.B. Hoj, et al., Selective induction of mitochondrial chaperones in response to loss of the mitochondrial genome, *Eur. J. Biochem.* 240 (1) (1996) 98–103.
- [48] T. Yoneda, C. Benedetti, F. Urano, S.G. Clark, H.P. Harding, D. Ron, Compartment-specific perturbation of protein handling activates genes encoding mitochondrial chaperones, *J. Cell Sci.* 117 (Pt 18) (2004) 4055–4066.
- [49] R.B. Hamanaka, N.S. Chandel, Mitochondrial reactive oxygen species regulate cellular signaling and dictate biological outcomes, *Trends Biochem. Sci.* 35 (9) (2010) 505–513.
- [50] C.M. Haynes, K. Petrova, C. Benedetti, Y. Yang, D. Ron, ClpP mediates activation of a mitochondrial unfolded protein response in *C. elegans*, *Dev. Cell.* 13 (4) (2007) 467–480.
- [51] C.M. Haynes, D. Ron, The mitochondrial UPR – protecting organelle protein homeostasis, *J. Cell Sci.* 123 (Pt 22) (2010) 3849–3855.
- [52] T. Iwawaki, R. Akai, K. Kohno, M. Miura, A transgenic mouse model for monitoring endoplasmic reticulum stress, *Nat. Med.* 10 (1) (2004) 98–102.
- [53] T. Iwawaki, R. Akai, S. Yamanaka, K. Kohno, Function of IRE1 $\alpha$  in the placenta is essential for placental development and embryonic viability, *Proc. Natl. Acad. Sci. U. S. A.* 106 (39) (2009) 16657–16662.
- [54] J. Heijmans, J.F. van Lidth de Jude, B.K. Koo, S.L. Rosekrans, M.C. Wielenga, M. van de Wetering, et al., ER stress causes rapid loss of intestinal epithelial stemness through activation of the unfolded protein response, *Cell Rep.* 3 (4) (2013) 1128–1139.
- [55] J.A. Barbour, N. Turner, Mitochondrial stress signaling promotes cellular adaptations, *Int. J. Cell Biol.* 2014 (2014) 156020.
- [56] Y. Ma, Y. Shimizu, M.J. Mann, Y. Jin, L.M. Hendershot, Plasma cell differentiation initiates a limited ER stress response by specifically suppressing the PERK-dependent branch of the unfolded protein response, *Cell Stress Chaperones* 15 (3) (2010) 281–293.
- [57] H.W. Yung, M. Hemberger, E.D. Watson, C.E. Senner, C.P. Jones, R.J. Kaufman, et al., Endoplasmic reticulum stress disrupts placental morphogenesis: implications for human intrauterine growth restriction, *J. Pathol.* 228 (2012) 554–564.
- [58] D. Scheuner, B. Song, E. McEwen, C. Liu, R. Laybutt, P. Gillespie, et al., Translational control is required for the unfolded protein response and in vivo glucose homeostasis, *Mol. Cell.* 7 (6) (2001) 1165–1176.
- [59] T. Kawakami, M. Yoshimi, Y. Kadota, M. Inoue, M. Sato, S. Suzuki, Prolonged endoplasmic reticulum stress alters placental morphology and causes low birth weight, *Toxicol. Appl. Pharmacol.* 275 (2) (2014) 134–144.
- [60] R. Delsite, S. Kachhap, R. Anbazhagan, E. Gabrielson, K.K. Singh, Nuclear genes involved in mitochondria-to-nucleus communication in breast cancer cells, *Mol. Cancer* 1 (2002) 6.
- [61] S. Gispert, D. Parganlija, M. Klinckenberg, S. Drose, I. Wittig, M. Mittelbronn, et al., Loss of mitochondrial peptidase Clpp leads to infertility, hearing loss plus growth retardation via accumulation of CLPX, mtDNA and inflammatory factors, *Hum. Mol. Genet.* 22 (24) (2013) 4871–4887.
- [62] E. Jauniaux, J. Hempstock, N. Greenwold, G.J. Burton, Trophoblastic oxidative stress in relation to temporal and regional differences in maternal placental blood flow in normal and abnormal early pregnancies, *Am. J. Pathol.* 162 (2003) 115–125.
- [63] G.J. Burton, E. Jauniaux, D.S. Charnock-Jones, The influence of the intrauterine environment on human placental development, *Int. J. Dev. Biol.* 54 (2010) 303–312.
- [64] H.W. Yung, M. Cox, M. Tissot van Patot, G.J. Burton, Evidence of endoplasmic reticulum stress and protein synthesis inhibition in the placenta of non-native women at high altitude, *FASEB J.* 26 (2012) 1970–1981.
- [65] F. Colleoni, N. Padmanabhan, H.W. Yung, E.D. Watson, I. Cetin, M. Tissot van Patot, et al., Suppression of mitochondrial electron transport chain function in the hypoxic human placenta: a role for miR-210 and protein synthesis inhibition, *PLoS One* 8 (2013) e55194.
- [66] M.C. Tissot van Patot, M. Valdez, V. Becky, T. Cindrova-Davies, J. Johns, L. Zwerdling, et al., Impact of pregnancy at high altitude on placental morphology in non-native women with and without preeclampsia, *Placenta* 30 (6) (2009) 523–528.
- [67] H. Matheson, J.H. Veerbeek, D.S. Charnock-Jones, G.J. Burton, H.W. Yung, Morphological and molecular changes in the murine placenta exposed to normobaric hypoxia throughout pregnancy, *J. Physiol.* 594 (5) (2016) 1371–1388.
- [68] M. Fahling, Surviving hypoxia by modulation of mRNA translation rate, *J. Cell Mol. Med.* 13 (9A) (2009) 2770–2779.
- [69] M.C. Tissot van Patot, A.J. Murray, V. Beckey, T. Cindrova-Davies, J. Johns, L. Zwerdling, et al., Human placental metabolic adaptation to chronic hypoxia, high altitude: hypoxic preconditioning, *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 298 (1) (2010) R166–R172.
- [70] G.L. Semenza, P.H. Roth, H.M. Fang, G.L. Wang, Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1, *J. Biol. Chem.* 269 (38) (1994) 23757–23763.
- [71] J.W. Kim, I. Tchernyshyov, G.L. Semenza, C.V. Dang, HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia, *Cell Metab.* 3 (3) (2006) 177–185.
- [72] A.J. Murray, Oxygen delivery and fetal-placental growth: beyond a question of supply and demand? *Placenta* 33 (Suppl 2) (2012) e16–22.
- [73] D.Z. Levett, E.J. Radford, D.A. Menassa, E.F. Graber, A.J. Morash, H. Hoppeler, et al., Acclimatization of skeletal muscle mitochondria to high-altitude hypoxia during an ascent of Everest, *FASEB J.* 26 (4) (2012) 1431–1441.
- [74] D.Z. Levett, A. Vigano, D. Capitanio, M. Vasso, S. De Palma, M. Moriggi, et al., Changes in muscle proteomics in the course of the caudwell research expedition to Mt. Everest, *Proteomics* 15 (1) (2015) 160–171.
- [75] L.M. Edwards, A.J. Murray, D.J. Tyler, G.J. Kemp, C.J. Holloway, P.A. Robbins, et al., The effect of high-altitude on human skeletal muscle energetics: P-MRS results from the Caudwell Xtreme Everest expedition, *PLoS One* 5 (5) (2010) e10681.
- [76] I. Brosens, R. Pijnenborg, L. Vercruyse, R. Romero, The “Great Obstetrical Syndromes” are associated with disorders of deep placentation, *Am. J. Obstet. Gynecol.* 204 (3) (2011) 193–201.
- [77] H.W. Yung, D. Atkinson, T. Campion-Smith, M. Olovsson, D.S. Charnock-Jones, G.J. Burton, Differential Activation of Placental Unfolded Protein Response Pathways Implies Heterogeneity in Causation of Early- and Late-onset Pre-eclampsia, *J. Pathol.* 234 (2014) 262–276.
- [78] K. Wataba, T. Saito, M. Takeuchi, M. Nakayama, N. Suehara, R. Kudo, Changed expression of heat shock proteins in various pathological findings in placentas with intrauterine fetal growth restriction, *Med. Electron Microsc.* 37 (3) (2004) 170–176.
- [79] T.-H. Hung, D.S. Charnock-Jones, J.N. Skepper, G.J. Burton, Secretion of tumour necrosis factor- $\alpha$  from human human placental tissues induced by hypoxia-reoxygenation causes endothelial cell activation in vitro: a potential mediator of the inflammatory response in preeclampsia, *Am. J. Pathol.* 164 (2004) 1049–1061.
- [80] T. Cindrova-Davies, H.W. Yung, J. Johns, O. Spasic-Boskovic, S. Korolchuk, E. Jauniaux, et al., Oxidative stress, gene expression, and protein changes induced in the human placenta during labor, *Am. J. Pathol.* 171 (2007) 1168–1179.
- [81] T. Cindrova-Davies, From placental oxidative stress to maternal endothelial dysfunction, *Placenta Suppl. A* 30 (2009) S55–S65.
- [82] T.H. Hung, S.F. Chen, T.T. Hsieh, L.M. Lo, M.J. Li, Y.L. Yeh, The associations between labor and delivery mode and maternal and placental oxidative stress, *Reprod. Toxicol.* 31 (2) (2011) 144–150.
- [83] T. Cindrova-Davies, O. Spasic-Boskovic, E. Jauniaux, D.S. Charnock-Jones, G.J. Burton, Nuclear factor- $\kappa$ B, p38, and stress-activated protein kinase mitogen-activated protein kinase signaling pathways regulate proinflammatory cytokines and apoptosis in human placental explants in response to oxidative stress: effects of antioxidant vitamins, *Am. J. Pathol.* 170 (5) (2007) 1511–1520.
- [84] C. Xu, B. Bailly-Maitre, J.C. Reed, Endoplasmic reticulum stress: cell life and death decisions, *J. Clin. Invest.* 115 (10) (2005) 2656–2664.
- [85] K. Zhang, R.J. Kaufman, From endoplasmic-reticulum stress to the inflammatory response, *Nature* 454 (7203) (2008) 455–462.
- [86] J. Deng, P.D. Lu, Y. Zhang, D. Scheuner, R.J. Kaufman, N. Sonenberg, et al., Translational repression mediates activation of nuclear factor kappa B by phosphorylated translation initiation factor 2, *Mol. Cell Biol.* 24 (23) (2004) 10161–10168.
- [87] T.Y. Khong, F. De Wolf, W.B. Robertson, I. Brosens, Inadequate maternal vascular response to placentation in pregnancies complicated by preeclampsia and by small-for-gestational age infants, *Br. J. Obstet. Gynaecol.* 93 (10) (1986) 1049–1059.
- [88] S. Sohlberg, A. Mulic-Lutvica, P. Lindgren, F. Ortiz-Nieto, A.K. Wikstrom, J. Wikstrom, Placental perfusion in normal pregnancy and early and late preeclampsia: a magnetic resonance imaging study, *Placenta* 35 (3) (2014) 202–206.
- [89] S. Sohlberg, A.K. Wikstrom, M. Olovsson, P. Lindgren, O. Axelsson, A. Mulic-Lutvica, et al., In vivo (31)P-MR spectroscopy in normal pregnancy, early and late preeclampsia: a study of placental metabolism, *Placenta* 35 (5) (2014) 318–323.
- [90] Y. Ovadya, V. Krizhanovskiy, Senescent cells: SASPected drivers of age-related pathologies, *Biogerontology* 15 (6) (2014) 627–642.
- [91] C. Correia-Melo, J.F. Passos, Mitochondria: are they causal players in cellular senescence? *Biochim. Biophys. Acta* 1847 (11) (2015) 1373–1379.
- [92] Y.S. Yoon, H.O. Byun, H. Cho, B.K. Kim, G. Yoon, Complex II defect via down-

- regulation of iron-sulfur subunit induces mitochondrial dysfunction and cell cycle delay in iron chelation-induced senescence-associated growth arrest, *J. Biol. Chem.* 278 (51) (2003) 51577–51586.
- [93] P. Stockl, E. Hutter, W. Zwerschke, P. Jansen-Durr, Sustained inhibition of oxidative phosphorylation impairs cell proliferation and induces premature senescence in human fibroblasts, *Exp. Gerontol.* 41 (7) (2006) 674–682.
- [94] O. Moiseeva, V. Bourdeau, A. Roux, X. Deschenes-Simard, G. Ferbeyre, Mitochondrial dysfunction contributes to oncogene-induced senescence, *Mol. Cell Biol.* 29 (16) (2009) 4495–4507.
- [95] A. Chuprin, H. Gal, T. Biron-Shental, A. Biran, A. Amiel, S. Rozenblatt, et al., Cell fusion induced by ERVWE1 or measles virus causes cellular senescence, *Genes. Dev.* 27 (21) (2013) 2356–2366.
- [96] D. Goldman-Wohl, S. Yagel, United we stand not dividing: the syncytiotrophoblast and cell senescence, *Placenta* 35 (6) (2014) 341–344.
- [97] B.J. Martin, S.S. Spicer, Ultrastructural features of cellular maturation and aging in human trophoblast, *J. Ultrastruct. Res.* 43 (1973) 133–149.
- [98] C.J. Jones, H. Fox, Ultrastructure of the placenta in prolonged pregnancy, *J. Pathol.* 126 (3) (1978) 173–179.
- [99] N.M. Fogarty, A.C. Ferguson-Smith, G.J. Burton, Syncytial knots (tenney-parker changes) in the human placenta: evidence of loss of transcriptional activity and oxidative damage, *Am. J. Pathol.* 183 (1) (2013) 144–152.
- [100] G.J. Burton, E. Jauniaux, Placental oxidative stress; from miscarriage to preeclampsia, *J. Soc. Gynecol. Investig.* 11 (2004) 342–352.
- [101] M.L. Tjoa, T. Cindrova-Davies, O. Spasic-Boskovic, D.W. Bianchi, G.J. Burton, Trophoblastic oxidative stress and the release of cell-free feto-placental DNA, *Am. J. Pathol.* 169 (2006) 400–404.
- [102] Y.M. Lo, M.S. Tein, T.K. Lau, C.J. Haines, T.N. Leung, P.M. Poon, et al., Quantitative analysis of fetal DNA in maternal plasma and serum: implications for noninvasive prenatal diagnosis, *Am. J. Hum. Genet.* 62 (4) (1998) 768–775.
- [103] C. Burdon, C. Mann, T. Cindrova-Davies, A.C. Ferguson-Smith, G.J. Burton, Oxidative stress and the induction of cyclooxygenase enzymes and apoptosis in the murine placenta, *Placenta* 28 (2007) 724–733.
- [104] C.E. Aiken, T. Cindrova-Davies, M.H. Johnson, Variations in mouse mitochondrial DNA copy number from fertilization to birth are associated with oxidative stress, *Reprod. Biomed. Online* 17 (6) (2008) 806–813.
- [105] R.J. Levine, S.E. Maynard, C. Qian, K.H. Lim, L.J. England, K.F. Yu, et al., Circulating angiogenic factors and the risk of preeclampsia, *N. Engl. J. Med.* 350 (2004) 672–683.
- [106] M. Mizuuchi, T. Cindrova-Davies, M. Olovsson, D.S. Charnock-Jones, G.J. Burton, H.W. Yung, Placental endoplasmic reticulum stress negatively regulates transcription of placental growth factor via ATF4 and ATF6beta: implications for the pathophysiology of human pregnancy complications, *J. Pathol.* 238 (2016) 550–561.
- [107] P. Davy, M. Nagata, P. Bullard, N.S. Fogelson, R. Allsopp, Fetal growth restriction is associated with accelerated telomere shortening and increased expression of cell senescence markers in the placenta, *Placenta* 30 (6) (2009) 539–542.
- [108] T. Biron-Shental, R. Sukenik-Halevy, Y. Sharon, L. Goldberg-Bittman, D. Kidron, M.D. Fejgin, et al., Short telomeres may play a role in placental dysfunction in preeclampsia and intrauterine growth restriction, *Am. J. Obstet. Gynecol.* 202(4) (381) (2010) e1–7.
- [109] T. Biron-Shental, D. Sadeh-Mestechkin, A. Amiel, Telomere homeostasis in IUGR placentas – A review, *Placenta* 39 (2016) 21–23.
- [110] A.R. Davalos, J.P. Coppe, J. Campisi, P.Y. Desprez, Senescent cells as a source of inflammatory factors for tumor progression, *Cancer Metastasis Rev.* 29 (2) (2010) 273–283.
- [111] L. Zhu, Z. Zhang, L. Zhang, Y. Shi, J. Qi, A. Chang, et al., HMGB1-RAGE signaling pathway in severe preeclampsia, *Placenta* 36 (10) (2015) 1148–1152.
- [112] D. Narendra, A. Tanaka, D.F. Suen, R.J. Youle, Parkin is recruited selectively to impaired mitochondria and promotes their autophagy, *J. Cell Biol.* 183 (5) (2008) 795–803.
- [113] D.F. Suen, D.P. Narendra, A. Tanaka, G. Manfredi, R.J. Youle, Parkin overexpression selects against a deleterious mtDNA mutation in heteroplasmic cybrid cells, *Proc. Natl. Acad. Sci. U. S. A.* 107 (26) (2010) 11835–11840.
- [114] S. Deegan, S. Saveljeva, A.M. Gorman, A. Samali, Stress-induced self-cannibalism: on the regulation of autophagy by endoplasmic reticulum stress, *Cellular and molecular life sciences, CMLS* 70 (14) (2013) 2425–2441.
- [115] S. Deegan, I. Koryga, S.A. Glynn, S. Gupta, A.M. Gorman, A. Samali, A close connection between the PERK and IRE arms of the UPR and the transcriptional regulation of autophagy, *Biochem. Biophys. Res. Commun.* 456 (1) (2015) 305–311.
- [116] T.H. Hung, S.F. Chen, L.M. Lo, M.J. Li, Y.L. Yeh, T.T. Hsieh, Increased autophagy in placentas of intrauterine growth-restricted pregnancies, *PLoS One* 7 (7) (2012) e40957.
- [117] S. Curtis, C.J. Jones, A. Garrod, C.H. Hulme, A.E. Heazell, Identification of autophagic vacuoles and regulators of autophagy in villous trophoblast from normal term pregnancies and in fetal growth restriction, *J. Matern. Fetal Neonatal Med.* 26 (4) (2013) 339–346.
- [118] S.Y. Oh, S.J. Choi, K.H. Kim, E.Y. Cho, J.H. Kim, C.R. Roh, Autophagy-related proteins, LC3 and Beclin-1, in placentas from pregnancies complicated by preeclampsia, *Reprod. Sci.* 15 (9) (2008) 912–920.
- [119] R. Akaiishi, T. Yamada, K. Nakabayashi, H. Nishihara, I. Furuta, T. Kojima, et al., Autophagy in the placenta of women with hypertensive disorders in pregnancy, *Placenta* 35 (12) (2014) 974–980.
- [120] B. Chen, M.S. Longtine, D.M. Nelson, Hypoxia induces autophagy in primary human trophoblasts, *Endocrinology* 153 (10) (2012) 4946–4954.
- [121] T.H. Hung, T.T. Hsieh, S.F. Chen, M.J. Li, Y.L. Yeh, Autophagy in the human placenta throughout gestation, *PLoS One* 8 (12) (2013) e83475.
- [122] S. Saito, A. Nakashima, Review: The role of autophagy in extravillous trophoblast function under hypoxia, *Placenta* 34 (Suppl.:S79–84) (2013).
- [123] K.D. Broad, E.B. Keverne, Placental protection of the fetal brain during short-term food deprivation, *Proc. Natl. Acad. Sci. U. S. A.* 108 (37) (2011) 15237–15241.
- [124] M.S. Longtine, A. Barton, B. Chen, D.M. Nelson, Live-cell imaging shows apoptosis initiates locally and propagates as a wave throughout syncytiotrophoblasts in primary cultures of human placental villous trophoblasts, *Placenta* 33 (12) (2012) 971–976.
- [125] M.S. Longtine, B. Chen, A.O. Odibo, Y. Zhong, D.M. Nelson, Caspase-mediated apoptosis of trophoblasts in term human placental villi is restricted to cytotrophoblasts and absent from the multinucleated syncytiotrophoblast, *Reproduction* 143 (1) (2012) 107–121.
- [126] G.J. Burton, J.N. Skepper, J. Hempstock, T. Cindrova, C.J.P. Jones, E. Jauniaux, A reappraisal of the contrasting morphological appearances of villous cytotrophoblast cells during early human pregnancy; evidence for both apoptosis and primary necrosis, *Placenta* 24 (2003) 297–305.
- [127] J. Hempstock, E. Jauniaux, N. Greenwold, G.J. Burton, The contribution of placental oxidative stress to early pregnancy failure, *Hum. Pathol.* 34 (2003) 1265–1275.
- [128] M.S. Longtine, B. Chen, A.O. Odibo, Y. Zhong, D.M. Nelson, Villous trophoblast apoptosis is elevated and restricted to cytotrophoblasts in pregnancies complicated by preeclampsia, IUGR, or preeclampsia with IUGR, *Placenta* 33 (2012) 352–359.
- [129] R.A. Simpson, T.M. Mayhew, P.R. Barnes, From 13 weeks to term, the trophoblast of human placenta grows by the continuous recruitment of new proliferative units: a study of nuclear number using the disector, *Placenta* 13 (1992) 501–512.
- [130] I.A. Lian, M. Loset, S.B. Mundal, M.H. Fenstad, M.P. Johnson, I.P. Eide, et al., Increased endoplasmic reticulum stress in decidual tissue from pregnancies complicated by fetal growth restriction with and without pre-eclampsia, *Placenta* 32 (11) (2011) 823–829.
- [131] H.J. Gao, Y.M. Zhu, W.H. He, A.X. Liu, M.Y. Dong, M. Jin, et al., Endoplasmic reticulum stress induced by oxidative stress in decidual cells: a possible mechanism of early pregnancy loss, *Mol. Biol. Rep.* 39 (9) (2012) 9179–9186.
- [132] A. Moffett, S.E. Hiby, A.M. Sharkey, The role of the maternal immune system in the regulation of human birthweight, *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 370 (1663) (2015).
- [133] L. Ulianic, G. Terrazzano, M. Annunziata, G. Ruggiero, F. Beguinot, B. Di Jeso, ER stress impairs MHC Class I surface expression and increases susceptibility of thyroid cells to NK-mediated cytotoxicity, *Biochim. Biophys. Acta* 1812 (4) (2011) 431–438.